

INTERVENTIONS FOR EMERGENCY SITUATIONS

INTERVENTIONS FOR LOW BLOOD SUGAR

Hypoglycemia (Low blood sugar) – Insulin Reaction (**Must be accompanied to Health Office**):

(Any blood sugar below _____ constitutes a low blood sugar.

If blood sugar cannot be obtained, treat based on symptoms? Yes _____ No _____

Mild reaction signs person might exhibit are:

hunger irritability shakiness sleepiness sweating pallor
other: _____

Person usually recognizes the symptoms? _____ No _____

Yes _____

Time reactions most frequently occur? _____

√ **Treat mild** low blood sugar as follows:

_____ glucose tabs _____ cup juice _____ cup regular pop

other: _____

Follow initial treatment with a snack of _____ in _____ minutes.

Moderate reaction signs person might exhibit are:

confusion slurred speech disoriented sleepiness change in personality
other: _____

√ **Treat moderate low** blood sugar as follows:

_____ tube glucose gel _____ tube cake decorating gel _____ cup juice

other: _____

Person should _____ should not _____ follow with a snack of _____ in _____ minutes
or once symptoms subside.

Severe reaction signs person might exhibit are:

unconscious episode seizure unable/unwilling to take gel or juice

√ For **severe low** blood sugar, **treat** as follows:

_____ cc. Glucagon injection (_____ units) Call 911 Notify parents

Order for Glucagon on file? Yes _____ No _____

Call parent in the event _____

INTERVENTIONS FOR HIGH BLOOD SUGAR:

Hyperglucemia:

A blood sugar _____ will _____ may _____ require an insulin administration (see insulin
above _____ dosages).

If blood sugar is greater than _____, test blood ketones _____ urine ketones _____.

Child will _____ will not _____ need supervision in testing ketones.

Notify parents if blood glucose is _____ or when ketones
above _____ are _____

Comments: _____
